

Calculation of the Electric Dipole Field

From the first antisymmetry law:

$$\underline{E} = -\underline{\nabla} \phi + \underline{\omega} \phi = -\frac{\partial A}{\partial t} - \omega_0 \underline{A} \quad (1)$$

and assuming that

$$\frac{\partial A}{\partial t} = 0 \quad (2)$$

then

$$\underline{E} = -\omega_0 \underline{A} \quad (3)$$

Assuming that

$$\omega_0 = \frac{mc^2}{2\pi\hbar} \quad (4)$$

then

$$\underline{E} = -\frac{mc^2}{2\pi\hbar} \underline{A} \quad (5)$$

Here $mc^2/(2\pi\hbar)$ is a universal constant. In general, the electric dipole field is:

$$\underline{E}(\underline{r}) = \frac{3\underline{n}(\underline{p} \cdot \underline{n}) - \underline{p}}{4\pi\epsilon_0 |\underline{r} - \underline{r}_0|^3} \quad (6)$$

Here \underline{n} is a unit vector from \underline{r} to \underline{r}_0 . If the electric pole moment is in the z axis then:

$$\underline{E} = E_r \underline{e}_r + E_\theta \underline{e}_\theta \quad (7)$$

$$= \frac{p}{4\pi\epsilon_0 r^3} (2\cos\theta \underline{e}_r + \sin\theta \underline{e}_\theta)$$

spherical polar coordinates.

Therefore:

$$\underline{A} = -\frac{p}{4\pi\epsilon_0 \omega_0 r^3} (2\cos\theta \underline{e}_r + \sin\theta \underline{e}_\theta) \quad (8)$$

In the general case:

$$\underline{A}(\underline{r}) = -\frac{1}{\omega_0} \left(\frac{3\underline{n}(\underline{p} \cdot \underline{n}) - \underline{p}}{4\pi\epsilon_0 |\underline{r} - \underline{r}_0|^3} \right) - (9)$$

This is the electric vector potential for a dipole field.

Note that

$$\underline{\nabla} \omega_0 = \underline{0} - (10)$$

and

$$\underline{\nabla} \times \underline{A} = \underline{0} - (11)$$

so: $\underline{\nabla} \times (\omega_0 \underline{A}) = \omega_0 \underline{\nabla} \cdot \underline{A} + \underline{A} \cdot \underline{\nabla} \omega_0 = \underline{0} - (12)$

is required for eq. (1) and

$$\underline{\nabla} \times \underline{E} = \underline{0} - (13)$$

The Faraday law of induction means that:

$$\underline{\nabla} \times \underline{E} + \frac{\partial \underline{B}}{\partial t} = \underline{0} - (14)$$

so

$$\frac{\partial \underline{B}}{\partial t} = \underline{0} - (15)$$

if it is assumed that:

$$\underline{B} = \underline{\nabla} \times \underline{A} - \underline{\omega} \times \underline{A} = \underline{0} - (16)$$

then eq. (15) is true automatically.

The vector identity law, together with eq. (16), mean that the spin connections can be calculated by computer algebra as follows:

$$\begin{aligned}\frac{\partial A_z}{\partial y} &= \omega_y A_z, & \frac{\partial A_y}{\partial z} &= -\omega_z A_y, \\ \frac{\partial A_x}{\partial z} &= \omega_z A_x, & \frac{\partial A_z}{\partial x} &= \omega_x A_z, \\ \frac{\partial A_y}{\partial x} &= \omega_x A_y, & \frac{\partial A_x}{\partial y} &= \omega_y A_x.\end{aligned}\quad - (17)$$

Therefore eqs. (1) and (8) must be expressed in Cartesian components. To avoid human error this is best done by computer algebra. A preliminary hand calculation gives (Note 385(2)):

$$\underline{A} = -\frac{\rho}{4\pi\epsilon_0\omega_0 r^5} \left(3xz \underline{i} + 3xy \underline{j} + (2z^2 - x^2 - y^2) \underline{k} \right).$$

where

$$r^5 = (x^2 + y^2 + z^2)^{5/2} \quad - (18)$$

$$r^5 = (x^2 + y^2 + z^2)^{5/2} \quad - (19)$$

The spic connections calculated in this way preserve symmetry. The spic connections have an interesting symptical structure.

The calculation can be repeated for the

$$\omega_0 = -\frac{c}{r} \quad - (20)$$